



© Hülya Sungurtekin,
© Fatih Sargın,
© Metin Kılınc,
© Mithat Kahramanoğlu,
© Mert Akbulut,
© Ferruh Ayoğlu,
© Ahmet Çalışkan,
© Simay Karaduman

Usefulness of APACHE II, SOFA, ISARIC/WHO 4C Mortality Score and CO-RADS for Mortality Prediction of Critically Ill Coronavirus Disease 2019 (Covid-19) Patients

APACHE II, SOFA, ISARIC/WHO 4C Mortalite Skoru ve CO-RADS'in Kritik Coronavirüs 2019 (Covid-19) Hastalarının Mortalite Tahmininde Kullanımı

Received/Geliş Tarihi : 19.04.2022
Accepted/Kabul Tarihi : 26.08.2022

Hülya Sungurtekin, Fatih Sargın, Metin Kılınc, Mithat Kahramanoğlu, Mert Akbulut, Simay Karaduman
Pamukkale University Faculty of Medicine,
Department of Anesthesiology and Reanimation,
Turkey

Ferruh Ayoğlu
Zonguldak Bülent Ecevit University Faculty of
Medicine, Department of Public Health, Zonguldak,
Turkey

Ahmet Çalışkan
Pamukkale University Faculty of Medicine,
Department of Medical Microbiology, Denizli, Turkey

Hülya Sungurtekin Prof. MD (✉),
Pamukkale University Faculty of Medicine,
Department of Anesthesiology and Reanimation,
Turkey

E-mail : hsungurtekin@yahoo.com

Phone : +90 532 789 43 30

ORCID ID : orcid.org/0000-0002-9453-5625

ABSTRACT Objective: It was aimed to report the Acute Physiology and Chronic Health Evaluation II (APACHE II) score, Sequential Organ Failure Assessment (SOFA) score, Glasgow Coma Scale (GCS), 4C mortality score and the COVID-19 Reporting and Data System (CO-RADS) in predicting the outcome of critically ill COVID-19 patients.

Materials and Methods: Patients with laboratory-confirmed COVID-19 infection or clinical and radiological confirmed COVID-19 infection who were admitted to adult ICU were included. Clinical characteristics, outcomes, APACHE II score, SOFA score, ISARIC/WHO 4C mortality score and CO-RADS classification were reported at admission.

Results: 276 patients were included in this study. The mean age was higher in non-survivor patients. The most common cause of hospitalization was respiratory failure (67%). The common co-morbidities were hypertension (51.8%), cardiac disease (43.4%) and diabetes (33.6%). Organ failure was present in 61.5% of the patients. The mean APACHE II, SOFA, GCS and 4C mortality scores were higher in non-survivor patients. 4C mortality and SOFA scores showed higher predictive accuracy for mortality with an area AUC 0.736 and 0.706, respectively. 4C mortality had sensitivity of 78.9% and specificity of 58.1% whereas of SOFA had a sensitivity of 78.9% and a specificity of 53.3%.

Conclusion: 4C mortality and SOFA scores could be a predictors of mortality in COVID-19 patients in the ICU.

Keywords: COVID-19, Intensive Care, CO-RADS classification, 4C Mortality score, SOFA score, and APACHE II score

ÖZ Amaç: Kritik Covid-19 hastalarının Mortalite tahmininde Akut Fizyoloji ve Kronik Sağlık Değerlendirmesi II (APACHE II) skoru, Sıralı Organ Yetmezliği Değerlendirmesi (SOFA) skoru, Glasgow Koma Skalası (GCS), 4C Mortalite skoru ve Covid-19 Raporlama ve Veri Sistemi (CO-RADS)'nin araştırılması amaçlanmıştır.

Gereç ve Yöntem: Laboratuvarca doğrulanmış Covid-19 enfeksiyonu veya klinik ve radyolojik olarak doğrulanmış Covid-19 enfeksiyonu olan erişkin yoğun bakım ünitesine kabul edilen hastalar dahil edildi. Klinik özellikler, sonuçlar, APACHE II skoru, SOFA skoru, ISARIC/WHO 4C mortalite skoru ve CO-RADS sınıflandırması yatış esnasında kaydedildi.

Bulgular: Bu çalışmaya 276 hasta dahil edildi. Ölen hastalarda yaş ortalaması daha yüksekti. En sık hastaneye yatış nedeni solunum yetmezliğiydi (%67). En sık eşlik eden hastalıklar hipertansiyon (%51,8), kalp hastalığı (%43,4) ve diyabet (%33,6) idi. Hastaların %61,5'unda organ yetmezliği mevcuttu. Ortalama APACHE II, SOFA, GCS ve 4C mortalite skorları ölen hastalarda daha yüksekti. Mortalite için 4C mortalite ve SOFA skorları, sırasıyla AUC 0.736 ve 0.706 alan ile daha yüksek öngörü bulundu. 4C mortalite skoru %78.9 duyarlılık ve %58.1 özgüllüğe sahipken, SOFA'nın duyarlılığı %78.9 ve özgüllüğü %53.3 idi.

Sonuç: 4C mortalite ve SOFA skorları, yoğun bakımdaki Covid-19 hastalarında mortalitenin tahmin ettirici bir göstergesi olabilir.

Anahtar Kelimeler: Covid-19, Yoğun Bakım, CO-RADS sınıflandırması, 4C Mortalite skoru, SOFA skoru and APACHE II skoru

Introduction

The severe acute respiratory syndrome coronavirus disease 2019 (Covid-19) began in Wuhan, China, and has spread worldwide, infecting millions of people since than December 2019. The Coronavirus disease 2019 pandemic has caused an intense loss of human life worldwide and presents an extraordinary challenge to public health systems and the world economy.

Predicting the outcomes in intensive care patients is very important in terms of both guiding the treatment and preventing unnecessary treatments. Various laboratory tests, clinical findings or scoring systems are used to predict outcomes in intensive care patients. Research and large-scale vaccination campaigns are ongoing for effective treatment of Covid-19. Meanwhile, it is very important to predict in-hospital mortality during hospitalization for Covid-19.

The APACHE II scores was designed to calculate the severity of disease of intensive care unit (ICU) patients and to predict mortality. The APACHE II score is calculated based on body temperature, heart and respiratory rate, mean arterial pressure, pH. Range of APACHE II is 0 to 71. Increasing score is associated with an increasing risk of hospital death¹. The SOFA score was designed to assess the severity of organ dysfunction in critically ill septic patients. The original SOFA score was studied from a cohort of 1449 patients admitted to ICUs in sixteen countries². It was published that a high correlation between hospital mortality and the SOFA score in Covid-19 patients. They reported that SOFA score was a risk factor for death in Covid-19 patients³. However the discriminant accuracy of the SOFA score for mortality prediction in patients with Covid-19 pneumonia requiring mechanical ventilation was poor⁴.

In a review for prediction models for Covid-19 patients, they identified 107 prognostic models for patients with a diagnosis of Covid-19. The suggested use of these models was not visibly described. The most frequently used categories of prognostic factors (included at least 20 times for any outcome) included age, comorbidities, vital signs, image features, sex, lymphocyte count, and C reactive protein⁵. It was recommended the models by Knight and colleagues⁶ and Jehi and colleagues⁷ are good candidates for validation studies in other data for prediction models. Knight et al⁶ published the International Severe Acute Respiratory and emerging Infections Consortium (ISARIC) World Health

Organization (WHO) 4C Mortality Score for Covid-19. ISARIC/WHO 4C mortality score includes the biological and clinical variables, like breathing rate, peripheral oxygen saturation, age, sex, GCS, urea, CRP levels and number of comorbidities. The score ranges from zero to twenty-one points. A score of ≤ 3 had a 1% mortality risk compared with 62% mortality risk for those with a score of ≥ 15 .

Chest computerized tomography (CT) scans are used as a valuable tool in the diagnostic process of Covid-19 viral pneumonia cases. Chest CT specificity (82.9% to 96%) and sensitivity (80% to 90%) were reported to be higher than real-time reverse transcriptase-polymerase chain reaction (RT-PCR) testing for Covid-19 diagnosis. This highlights the need to recognize and understand imaging findings of the lungs⁸. Covid-19 Reporting and Data System (CO-RADS) is published in Mid-March of 2020 that grades the findings on how likely the diagnosis of Covid-19 is. It was evaluated using 105 randomly selected chest CT scans of patients admitted to the emergency department with clinical suspicion of Covid-19. CO-RADS system has seven categories. Categories zero and grade 1 to 6. Grade 1 to 6 means that from very low risk to proven infection with a positive RT-PCR assay. The system very well in estimating Covid-19 in patients with moderate to severe clinical disease⁹.

The objective of this paper is to report APACHE II score, SOFA score, the 4C mortality score and the CO-RADS classification in predicting outcome of Covid-19 in the ICU.

Materials and Methods

This is a single-center; retrospective cohort study that was analyzed anonymized data. Between September 1, 2020 and January 30, 2021 (the second wave of the Covid-19 pandemic), all adult patients (over than 18 years of the age) with RT-PCR assay confirmed Covid-19 infection or clinical and radiological confirmed Covid-19 infection were included.

Diagnosis of Covid-19 was accepted according to these findings: 1) Positive result of RT-PCR assay for Covid-19, 2), Typical Covid-19 lung CT scan abnormalities, 3) Covid-19 clinical findings and symptoms and/or the recent contact and/or travel history with certain case of Covid-19, associated with CO-RADS 3, 4 chest CT scan. Laboratory confirmation for Covid-19 was defined as a positive result of RT-PCR test from a specimen collected on an endotracheal aspirate or nasopharyngeal swab. RT-PCR assays (Covid-19 RT-qPCR, Bio-Speedy) were performed according to the protocol

approved by WHO in the General Office of Public Health Microbiology Reference Laboratory and laboratories in the specified areas¹⁰. Patient's data was obtained from hospital information systems. Data was recorded on daily basis.

Data Collection

The demographics and characteristics, clinical findings, therapies and laboratory data were recorded. The demographics and characteristics were age, sex, smoking history, APACHE II, GCS, SOFA, the 4C mortality score, laboratory data, clinical symptoms or signs, the recent exposure and travel history, comorbidities. The demographics and characteristics APACHE II, GCS, SOFA, the 4C mortality score, clinical symptoms or signs, the recent exposure and travel history, comorbidities were recorded at admission.

Laboratory data consisted of complete blood count (CBC), alkaline phosphatase, alanine aminotransferases (ALT), aspartate aminotransferases (AST), lactate dehydrogenase (LDH), serum creatinine (Cr), serum potassium, phosphate, sodium, D-dimer, prothrombin time (PT), international normalized ratio (INR), partial thromboplastin time (PTT), serum C-reactive protein (CRP), and serum procalcitonin. Invasive and noninvasive mechanical ventilation parameters were recorded. Arterial blood gas analysis were performed according to the patient's needs.

The ISARIC/WHO 4C mortality score was calculated on admission for each patient. This score includes eight parameters: age, sex, peripheral oxygen saturation, respiratory rate, number of comorbidities, level of consciousness (assessed using the GCS) and results of laboratory tests: serum urea and CRP⁶. Computerized tomography was performed at 1st day and as needed. All chest CT scans were performed without contrast agent and with a section thickness of 5 mm. The chest CT scans were reported according to CO-RADS. The CO-RADS scoring system has classification such as CO-RADS category 0 for technically insufficient imaging and CO-RADS category 6 for the confirmed disease through RT-PCR testing⁹.

Outcome

Patients' length of mechanical ventilation in ICU and discharge status (non-survivor, survivor) was entered patient's data form. Patients were treated according to the guide was prepared by scientific committee of the Ministry of Health and available literature¹¹. Therapies of the patients were documented daily. Since there was not enough evidence at the beginning of the pandemic, we did not

routinely use steroids in our patients. Microbial cultures from blood tracheal aspirate, and urine were taken at admission and in need of clinical situation.

Statistical Analysis

Statistical analyses were performed with SPSS for Windows 19 software. Continuous variable are shown as mean (standard deviation), median and minimum-maximum, categorical variables were reported as frequency with odds ratio and 95% confidence interval. The conformity of data to normal distribution were evaluated with Shapiro-Wilk test. The comparisons between groups were evaluated with Mann-Whitney U test for continuous variables, Chi-square test and Fisher's Exact Chi-square test for categorical values. Receiver operating curves (ROC) of APACHE II, SOFA, the 4C mortality score and CO-RADS were presented with area under curve (AUC), 95% confidence interval, cut-off value, sensitivity, specificity. Statistical analyses were evaluated with 95% confidence interval and $p < 0.05$ was accepted as a significant difference.

Results

Two hundred seventy six patients included to this study. Total 276 patients were grouped into two groups, depending on the survival status. One hundred seventy one patients (61.9%) were included in the non-survivor group. The rest of 105 patients (38.1%) had included in the survivor group. Total female and male patient were 98(36.6%) and 178(64.4%) respectively. The mean age was 68.6(13.3) years for all patients. In non-survivor patient group, the mean age were higher than in survivor patients ($P < .001$) (Table 1 and 2). One hundred eighty nine patients (68.4%) were admitted from emergency department. The most common cause of hospitalization was respiratory failure with a rate of 67%. The second most common cause was sepsis (33%). The reasons for hospitalization was statistically different between survivor and non-survivor patients. Sepsis is higher in non-survivor group than survivor group at admission ($P < .001$).

Two hundred twenty nine patients have comorbidities. Comorbidities have higher in the non-survivor group than survivor group. The most common co-morbidities were hypertension (51.8%), cardiac disease (43.4%) and diabetes (33.6%). Compared to the survivor group, non-survivor group had a significantly higher proportion of cardiac disease and pulmonary disease; the differences were statistically significant ($P < .05$). Dyspnea was the most common

symptom. Ninety five percent of the patients have dyspnea. The other common symptoms were, coughing and fever. Coughing was significantly higher in non-survivor patients than survivor patients were ($P=.003$) (Table 2).

Heart rate and FiO_2 was lower in survivor patients than non-survivor patients ($P<.001$), whereas the mean arterial pressure, was higher in survivor patients ($P =.003$). In

terms of comparison laboratory biomarkers taken during admission between 2 groups, it was found INR, D Dimer, creatinine, urea, CRP, AST, procalcitonin, lactate and IL 6 were significantly lower in the survivor group ($P<.001$) (Table 3). Chest radiography was available in 207 patients at admission. Pleural effusion were seen significantly higher in non-survivor patients ($P=.007$). Two hundred fifty two

Table 1. Demographics and clinical characteristics of the patients

Variable	Alive			Ex			All			p
	Mean \pm SD	Median	Min-Max	Mean \pm SD	Median	Min-Max	Mean \pm SD	Median	Min-Max	
Age	64.4 \pm 13.6	66	23-95	71.2 \pm 12.5	72	23-96	68.6 \pm 13.3	70	23-96	<0.001
BMI	29.7 \pm 3.4	30	22-42	29.5 \pm 3.6	29	20-46	29.6 \pm 3.5	29.3	20-46	0.566
Duration of symptom (day)	6.0 \pm 3.9	5	0-30	6.8 \pm 4.4	6	0-30	6.3 \pm 4.2	5	0-30	0.273
APACHE II	17.8 \pm 8.6	15	5-42	21.6 \pm 9.8	21	5-44	20.2 \pm 9.5	18	5-44	0.001
SOFA	5.0 \pm 2.5	4	0-12	7.4 \pm 3.5	7	0-18	6.5 \pm 3.3	6	0-18	<0.001
GCS	14.0 \pm 2.6	15	3-15	10.4 \pm 5.3	13	3-15	11.7 \pm 4.8	15	3-15	<0.001
4C Mortality	11.2 \pm 3.5	11	4-19	14.2 \pm 3.3	15	6-21	13.0 \pm 3.7	13	4-21	<0.001
CO-RADS	5.0 \pm 0.9	5	2-6	5.0 \pm 0.9	5	2-6	5.0 \pm 0.9	5	2-6	0.784
Fever	37.0 \pm 0.5	37	35.8-38.6	37.1 \pm 0.8	37	35.7-39.0	37.0 \pm 0.7	37	35.7-39.0	0.346
Heart rate (/min)	90.7 \pm 12.7	89	60-130	99.0 \pm 15.5	100	52-150	95.8 \pm 15.0	96	52-150	<0.001
Mean arterial pressure (mmHg)	81.1 \pm 9.1	80	66-120	77.7 \pm 12.7	75	50-122	79.0 \pm 11.5	78	50-122	0.003
FiO_2	50.9 \pm 7.8	50	40-80	63.1 \pm 10.1	60	40-90	58.5 \pm 11.0	60	40-90	<0.001
SPO_2	91.9 \pm 3.0	92	78-99	90.9 \pm 5.0	91	62-100	91.3 \pm 4.3	91	62-100	0.053
PEEP	6.0 \pm 1.1	6	5-10	7.8 \pm 1.9	8	5-14	7.1 \pm 1.9	7	5-14	<0.001
Tidal volume	388.6 \pm 46.2	390	320-450	418.3 \pm 43.9	420	300-600	416.0 \pm 44.6	410	300-600	0.032
Compliance mL cmH_2O	23.4 \pm 5.0	22	20-34.7	24.5 \pm 6.2	24	12-45	24.4 \pm 6.1	23.6	12-45	0.549
Driving Pressure	15.5 \pm 4.7	14	10-25	15.6 \pm 4.4	14	9-24	15.6 \pm 4.4	14	9-25	0.637
PaO_2/FiO_2	143.2 \pm 50.5	134	75-381	125.2 \pm 51.4	112	45-353	132.0 \pm 51.7	122	45-381	<0.001
Max PEEP during ICU	6.8 \pm 1.6	7	5-12	10.3 \pm 2.5	10	5-16	8.9 \pm 2.7	8	5-16	<0.001
Min PEEP during ICU	5.2 \pm 0.5	5	5-8	7.1 \pm 1.6	7	5-13	6.4 \pm 1.6	6	5-13	<0.001
Max PaO_2/FiO_2	279.3 \pm 74.8	261.5	132-520	186.7 \pm 70.4	180	17-456	221.7 \pm 89.7	220	17-520	<0.001
Min PaO_2/FiO_2	136.8 \pm 43.4	127	65-319	100.8 \pm 34.3	95	43-213	114.4 \pm 41.8	108	43-319	<0.001
Max PaO_2	109.9 \pm 34.9	99	55-202	98.8 \pm 37.6	88	42-212	103.0 \pm 36.9	93	42-212	0.003
Min PaO_2	61.6 \pm 22.0	54	41-185	59.1 \pm 19.7	52	30-132	60.1 \pm 20.6	53	30-185	0.204
Invasive mechanical ventilation (day)	9.6 \pm 8.4	7.5	1-27	7.1 \pm 7.7	4	1-36	7.2 \pm 7.7	4	1-36	0.197
Noninvasive mechanical ventilation (day)	7.9 \pm 5.7	6	1-26	4.8 \pm 4.4	3	1-27	6.6 \pm 5.4	5	1-27	<0.001

Table 2. Data frequencies of the patients

Variable	Alive (n=105)	Ex (n=171)	P	OR	CI	Variable	Alive (n=105)	Ex (n=171)	P	OR	CI
Gender (F/M)	36/69	62/109	0.740	0.92	0.51-1.53	CT (-/+)	8/97	16/155	0.782 ^a	0.80	0.33-1.94
Comorbidity (-/+)	27/78	20/151	0.004 ^a	2.61	1.38-4.95	CT Bilateral Ground Glass Opacification (-/+)	17/77	44/127	0.207 ^a	0.64	0.34-1.19
Hypertension (-/+)	58/47	75/96	0.066	1.58	0.97-2.58	CT Pleural Effusion (-/+)	80/14	119/52	0.008 ^a	2.50	1.30-4.81
Diabetes Mellitus (-/+)	66/39	114/57	0.519	0.85	0.51-1.41	CT Nodul (-/+)	86/6	163/8	0.712 ^b	0.70	0.24-2.09
Cardiac Disease (-/+)	70/35	86/85	0.008	1.98	1.19-3.27	Therapies and interventions					
Pulmonary Disease (-/+)	93/12	130/41	0.016	2.44	1.22-4.90	Recruitment (-/+)	103/2	146/25	0.001	8.82	2.04-38.050
Malignity (-/+)	93/12	138/33	0.121 ^a	1.85	0.91-3.77	Prone position (-/+)	69/36	156/15	<0.001	0.18	0.10-0.36
Chronic Renal Failure (-/+)	94/11	145/26	0.349 ^a	1.53	0.72-3.25	Renal replacement therapy (-/+)	104/1	158/13	0.031 ^a	8.56	1.10-66.39
Immune deficiency (-/+)	104/1	166/5	0.524 ^b	3.13	0.36-27.19	Plasma therapy (-/+)	99/6	158/13	0.721 ^a	1.36	0.45-3.69
Symptom (-/+)	3/102	1/170	0.311 ^b	5.00	0.51-48.70	Hidroksi Chloroquine (-/+)	99/6	167/4	0.262 ^a	0.40	0.11-1.44
Fever (-/+)	77/28	124/47	0.882	1.04	0.60-1.80	Antibiotic therapy (-/+)	5/100	6/165	0.825 ^b	1.38	0.41-4.62
Dyspnea (-/+)	6/99	8/163	0.922 ^a	1.24	0.42-3.66	Antiviral (-/+)	1/104	4/167	0.740 ^b	0.40	0.04-3.64
Cough (-/+)	45/60	105/66	0.003	0.47	0.29-0.77	Vasopressor (-/+)	93/12	17/154	<0.001	70.21	32.1-153.5
Smoking (-/+)	74/31	127/44	0.492	0.83	0.48-1.42	Tocilizumab (-/+)	98/7	162/9	0.827 ^a	0.78	0.28-2.16
Contact history (-/+)	80/25	136/35	0.513	0.82	0.46-1.48	Cytokine Adsorption (-/+)	104/1	157/14	0.021 ^a	9.27	1.20-71.59
						Aspirin (-/+)	0/105	2/169	0.383 ^b	0.54 [*]	0.06-5.21
Reason for admission (Respiratory failure/ Sepsis)	84/21	101/70	<0.001	2.77	1.57-4.89	Low molecular weight heparin(-/+)	0/105	2/169	0.383 ^b	0.54 [*]	0.06-5.21
PCR at admission (-/+)	12/93	5/166	0.009 ^a	4.28	1.46-12.53	Microbial culture (-/+)	74/31	116/55	0.646	1.13	0.67-1.92
Radiography (-/+)	27/78	42/129	0.830	1.06	0.61-1.86	Organ failure (-/+)	49/56	57/114	0.027	1.75	1.06-2.88
Radiography bilateral infiltrate (-/+)	29/76	44/127	0.730	1.10	0.64-1.91	Organ failure (-/ single/multiple)	49/52/4	57/55/59	<0.001	-	-
Radiography effusion (-/+)	92/13	125/46	0.007 ^a	2.60	1.33-5.10	Invasive mechanical ventilation (-/+)	95/10	4/167	<0.001	396.6	121.1-1299.0
Radiography ground glass opacities (-/+)	30/75	47/124	0.845	1.06	0.62-1.81	Noninvasive mechanical ventilation (-/+)	3/102	74/97	<0.001	0.04	0.01-0.13

^aYates corrected chi-square, ^bFisher's Exact chi-square

Table 3. Laboratory data of the patients

Variable	Alive			Ex			All			P
	Mean ± SD	Median	Min-Max	Mean ± SD	Median	Min-Max	Mean ± SD	Median	Min-Max	
White blood cell (per mm ³)	10893.1±5811.0	9720	100-28900	13116.3±13903.8	11090	340-171000	12270.5±11551.8	10870	100-171000	0.110
Lymphocyte (per mm ³)	822.7±1356.4	530	78-10100	684.0±459.3	520	100-2800	737.0±912.7	520	78-10100	0.883
Platelet (per mm ³)	1061.4±8369.7	225	36-86000	230.2±167.3	216	15-1900	546.4±5164.6	218	15-86000	0.232
Neutrophil to lymphocyte ratio	16.3±11.4	13.2	1.5-67	17.3±12.8	15	1.2-82	16.9±12.3	14.2	1.2-82	0.679
INR	1.2±0.5	1.1	0.9-4.6	1.4±1.8	1.2	0.9-9.3	1.3±0.7	1.2	0.9-9.3	<0.001
D-dimer ng/mL	1991.9±5589.1	570	48-46000	3556.6±6716.2	1290	106-55046	2961.4±6346.5	915	48-55046	<0.001
Fibrinogen mg/dL	583.1±246.5	586	99-1567	541.0±248.6	528	97-2000	557.0±248.2	550	97-2000	0.166
Creatinine mg/dL	1.2±1.3	0.9	0.4-9.7	1.7±1.3	1.2	0.3-8.9	1.5±1.3	1.1	0.3-9.7	<0.001
Aspartate aminotransferase	54.3±155.9	33	8-1612	143.9±530.1	38	8-6145	109.8±429.9	35.5	8-6145	0.009
Alanine aminotransferase	38.4±34.5	30	241	103.7±329.9	24	2-2864	78.9±262.1	26	2-2864	0.376
Ferritin ug/L	794.2±549.1	676.5	54-2000	852.7±627.0	663	32-3162	830.3±598.1	668.5	32-3162	0.745
C-reactive protein	98.9±70.6	88	0.9-388	136.1±143.8	106	6.7-1654	121.9±122.4	99.5	0.9-1654	0.006
Procalcitonin	1.6±6.8	0.2	0.03-66	5.3±13.0	0.7	0.01-80	3.9±11.2	0.4	0.01-80	<0.001
Lactate ug/L	1.7±0.4	1.8	0.8-3.1	2.4±1.9	2	0.8-16	2.2±1.7	1.9	0.8-16	0.005
IL6	58.5±76.4	32.5	3.4-397.8	256.3±701.1	63.5	12.1-3700	163.7±521.1	51.2	3.4-3700	0.001
Amylase	73.7±57.6	51	18-233	87.9±76.0	59	10-338	80.8±67.3	58.5	10-338	0.553
Lipase	41.8±42.5	28.3	10-232	79.0±134.8	42.5	4-726	60.1±100.4	32.6	4-726	0.259
Urea	56.8±32.4	49	13-159	93.3±60.2	75	19-353	79.4±54.3	62	13-353	<0.001

patient got CT scan at admission. Same as the radiography pleural effusion were shown significantly higher in non-survivor patients on CT scan ($P=0.008$) (Table 2).

Seventy percent of the patients were underwent invasive mechanical ventilation therapy. Invasive mechanical ventilation therapy was applied to the majority of patients who died whereas noninvasive mechanical ventilation therapy was applied to the majority of survivor patients. This difference was statistically significant ($P<0.001$) (Table 2). Positive-end-expiratory respiration (PEEP) and tidal volume values statistically were different between survivor and non-survivor groups. When we compared arterial blood gas analysis, it was found that non-survivor group had lower PaO₂ and PaO₂: FiO₂ ratio than the survivor group ($P<0.001$) (Table 1). Therapies and interventions were given in Table 2. Recruitment maneuver, prone position, renal replacement therapy, vasopressor usage and cytokine adsorption were higher in non-survivor patients and were statistically different from survivor patients ($P<0.001$). Organ failure was present in 61.5% of all patients. Multiple organ failure was more common in the non-survivor -group. The length of invasive and noninvasive mechanical ventilation were statistically significantly longer in survivor group ($P<0.001$) Table 2.

The mean APACHE II, SOFA, GCS and the 4C mortality scores were higher in non-survivor patients than in survivor patients ($P<0.001$) (Table 2). The mean CO-RADS classification value was not different between groups. Receiver-operating characteristic curves were computed to assess the accuracy of scores and CO-RADS in predicting mortality. 4C mortality and SOFA scores

showed higher predictive accuracy for mortality with an area AUC 0.736 and 0.706 respectively ($P < .001$) (Figure 1) (Table 4). The 4C mortality cut-off value of 11.5 had a sensitivity of 78.9% and specificity of 58.1%. The cut-off value of SOFA was 4.5, which corresponded to a sensitivity of 78.9% and a specificity of 53.3%. CO-RADS have not good results for predicting mortality with 5.5 cut-off value in ICU patients ($P = .802$).

Discussion

It was aimed to report APACHE II score, SOFA score, the 4C mortality score and the CO-RADS classification in predicting outcome of Covid-19 in the ICU in this study. Mortality was 61.9% and was higher in older patients and septic patients.

Previous studies from China, Europe, and United States (US), have described different mortality rates among critically ill patients ranging from 53.8% to 60.4% (12-14). Studies

from have reported that critically ill Covid-19 patients are generally older and have underlying medical conditions, such as hypertension and diabetes (12-14). Intensive Care National Audit & Research Centre (ICNARC) have evaluated data for 12,420 admissions of 10,873 patients critically ill with confirmed Covid-19. They reported that mortality was 55.9% patients with confirmed Covid-19 and any advanced respiratory support (13). In our study, we found also that older patients and patients have comorbidities higher mortality than others.

Covid-19 mainly in severe cases in addition to lung involves different organs such as heart, liver, and kidney, as well as hematological and nervous system, and induce multi-organ failure (15). In our study, patients have high percentage of respiratory failure and sepsis.

Despite advances in patient care, distance measures, and population vaccination campaigns Covid-19 still causes a high rate of death, especially in elderly and comorbid patients. As of 21 October 2021 more than 250 million cases of Covid-19 have been stated with more than 4 million deaths globally since than December 2019 (16). Hospitals all over the world are confronted with an influx of patients with Covid-19. There is an urgent need for a practical risk identification which will allow which patients are at the highest risk of death. Clinicians should consider prioritizing some therapies for patients at highest risk of clinical progression because of the optimize resource allocation and to guide management. Prediction models for Covid-19 are quickly entering the Covid-19 literature to support therapeutic choice making at a time when needed.

Yang et al (17) were reported that APACHE score and SOFA score were higher in died patients than in surviving patients. It was admitted fifty-two critically ill adult patients were with Covid-19 pneumonia in their study. Zou et al. (18) was aimed to investigate the APACHE score as a predictor for survival to facilitate decision-making for treatment in Wuhan.




Figure 1. Four score ROC

	Area under the curve (ROC)	95% CI	p	Cut-off	Sensitivity	Specificity
APACHE II	0.616	0.549-0.684	0.001	≥19.5	53.2	65.7
SOFA	0.706	0.644-0.768	<0.001	≥4.5	78.9	53.3
GCS	0.688	0.626-0.750	<0.001	≤13.5	50.9	83.8
CO-RADS	0.509	0.435-0.583	0.802	≥5.5	28.6	26.8
4C Mortality score	0.736	0.676-0.796	<0.001	≥11.5	78.9	58.1

In predicting hospital mortality, APACHE II score showed better discriminative ability (AUC, 0.966; 0.942–0.990, 95% CI) than SOFA score (AUC, 0.867; 0.808–0.926, 95% CI). APACHE II demonstrated a sensitivity and specificity of 96% and 86%, respectively in ICU patients. Stephens et al. (19) wrote a letter to the editor for Zou's study (18) and stated that raising questions about the calibration of APACHE II for Covid-19 patients. They noticed mortality higher than expected compared to relatively low APACHE II scores. One hundred and sixteen Covid-19 patients admitted to the ICU were retrospectively analyzed in Vandenbrande J et al study (20). They calculated APACHE II, APACHE IV scores and SOFA scores at admission. APACHE IV had a higher value for AUC than APACHE II (AUC 0.67 vs 0.63). APACHE IV and APACHE II have moderate discriminative power whereas the SOFA score had poor discriminative power (AUC:0.53) in all patients. In another paper, it was used multivariable logistic regression methods to investigate the risk factors associated with in-hospital mortality. It was reported increasing odds of in-hospital mortality associated with higher SOFA score in 191 adult Covid-19 patients (21). In our study, we also reported that higher APACHE II, SOFA, GCS and 4C Mortality scores in non-survivor patients. APACHE II and SOFA have AUC 0.616 and 0.706 respectively in our ICU patients. APACHE II and SOFA have same sensitivity such as 78.9%. SOFA score have better AUC data than APACHE II for predicting mortality in ICU Covid 19 patients.

The ISARIC/WHO 4C mortality score and the Jehi diagnostic model were reported as encouraging prediction models for Covid-19 (5). It was compared the ISARIC/WHO 4C mortality score to the CURB65, CRB65 and quick SOFA scores for to estimate 30-day mortality in patients with variety of respiratory infection in 606 patients. Fifty-three of 606 patients had Covid-19 infection. The ISARIC/WHO 4C mortality score had the highest AUR in Covid-19 patients for predicting mortality with value of 0.83 (22). Yildiz et al (23) prospectively evaluated 4C mortality score, Covid-GRAM, NEWS2, CURB-65 and compare them for progress of critical disease and poor outcome in a Covid-19 patients. The ISARIC/WHO 4C mortality score, Covid-GRAM, CURB-65 and NLR on admission showed strong predictive accuracy for mortality with an AUR curve of 0.80, 0.74, 0.74 and 0.76 respectively. 4C mortality score had the highest value for mortality prediction in Covid-19 hospitalized patients. In this model, a total score was calculated. In our study, the highest

predictive score was 4C mortality score in our critically ill patients.

Chest CT imaging shows a vital role in the diagnosis and evaluation of Covid-19 patients. The predictive value of high-resolution CT findings was reported in 181 mild-to-moderate and severe Covid-19 patients. It was shown that CT severity score might be a significant predictor of mortality in Covid-19 patients. The major chest CT finding is ground-glass opacity in both lungs, and multiple lobes in Covid-19 patients (24). The CO-RADS assessment scheme allows for the categorization of a chest CT scan. CO-RADS chest CT was greatly accurate for detecting Covid-19 pneumonia. The highest area under curve (AUC: 0.865) and accuracy (86.0%) was reported in the CO-RADS 4/5 group with a specificity of 112/132 (84.9%) and a sensitivity of 60/80 (88.2%) for diagnosing Covid-19 by RT-PCR test as the gold standard. Diagnostic value of CO-RADS on chest CT for the diagnosis of Covid-19 infection using the final clinical diagnosis as the standard of reference also have good result with area under curve 0.902) and accuracy (90.5%) in the CO-RADS 4/5 group (25). In our critically ill patients, CO-RADS was used for predicting of mortality in critically ill Covid 19 patients. CO-RADS was poor in predicting mortality in Covid 19 intensive care patients. In literature, there is no study that CO-RADS used Covid 19 for predicting mortality purposes.

If we were to comment on the mortality prediction of these five scorings used in our study, first, ROC studies may reach different values in different studies because of the difference in the patient population and the cut-off value chosen in the statistical method. In our study, CO-RADS did not give a significant result for predicting mortality. Although the most appropriate cut-off value is 5.5 and above, sensitivity and specificity values are quite low. The other four scores gave meaningful results and can be recommended for use mortality prediction. In our study, SOFA and 4C mortality score had the highest rates of accurate determination of mortality with 78.9%, but the probability of accurately determining survival in both was low.

This study has several limitations. It is a single center retrospective study and the study population was small. It was based on data mainly recorded hospital data base systems. Observer bias could be possible for CO-RADS evaluation.

Conclusion

SOFA and the 4C mortality score can be used to triage, guide decisions, and the clinical settings, to analyze early assessment of outcomes at admission. These scores may provide clinicians with a clue to discharge patients with low mortality scores or to manage early in patients whom need extra treatments. Future large studies should be aimed at developing and validating diagnostic and/or prognostic models for Covid-19 in ICU.

Ethics

Ethics Committee Approval: This is a single-center; retrospective study that was analyzed anonymized data from hospital database.

Informed Consent: Retrospective study.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: F.S., M.K., Mi.K., M.A., F.A., A.Ç., Concept: H.S., Design: H.S., Data Collection and Process: F.S., M.K., Mi.K., M.A., S.K., Analysis or Interpretation: H.S., F.A., Literature Search: H.S., F.S., M.K., Mi.K., M.A., Writing: H.S.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Crit Care Med.* 1985;13(10):818-829.
- Vincent JL, de Mendonça A, Cantraine F, Moreno R, Takala J, Suter PM, et al. Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on "sepsis-related problems" of the European Society of Intensive Care Medicine. *Crit Care Med.* 1998 Nov;26(11):1793-800.
- Yang Z, Hu Q, Huang F, Xiong S, Sun Y. The prognostic value of the SOFA score in patients with Covid-19: A retrospective, observational study. *Medicine (Baltimore).* 2021 Aug 13;100(32):e26900.
- Raschke RA, Agarwal S, Rangan P, Heise CW, Curry SC. Discriminant Accuracy of the SOFA Score for Determining the Probable Mortality of Patients With Covid-19 Pneumonia Requiring Mechanical Ventilation. *JAMA.* 2021;325(14):1469-1470.
- Wynants L, Van Calster B, Collins GS, Riley RD, Heinze G, Schuit E, et al. Prediction models for diagnosis and prognosis of Covid-19: systematic review and critical appraisal. *BMJ.* 2020 Apr 7;369:m1328. [published correction appears in *BMJ.* 2020 Jun 3;369:m2204].
- Knight SR, Ho A, Pius R, Buchan I, Carson G, Drake TM, et al. Risk stratification of patients admitted to hospital with Covid-19 using the ISARIC WHO Clinical Characterisation Protocol: development and validation of the 4C Mortality Score. *BMJ.* 2020 Sep 9;370:m3339.
- Jehi L, Ji X, Milinovich A, Erzurum S, Rubin BP, Gordon S, et al. Individualizing Risk Prediction for Positive Coronavirus Disease 2019 Testing: Results From 11,672 Patients. *Chest.* 2020 Oct;158(4):1364-1375.
- Majidi H, Niksolat F. Chest CT in patients suspected of Covid-19 infection: A reliable alternative for RT-PCR. *Am J Emerg Med.* 2020;38:2730-2732.
- Prokop M, van Everdingen W, van Rees Vellinga T, van Ufford JQ, Stöger L, Beenen L, et al. CO-RADS: A categorical CT assessment scheme for patients suspected of having Covid-19-definition and evaluation. *Radiology.* 2020;296:E97-104.
- World Health Organization. Coronavirus disease (Covid-19) technical guidance: laboratory testing for 2019-nCoV in humans (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/laboratory-guidance> accessed 14 Apr 2020).
- T. C. Ministry of Health general directorate of public health, Covid-19 (SARS-CoV-2 infection guide, Ankara. <https://Covid-19bilgi.saglik.gov.tr/depo/tedavi/Covid-19-EriskinHastaTedavisi.pdf> accessed 14 June 2020).
- Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ.* 2020 May 22;369:m1966.
- ICNARC report on Covid-19 in critical care: England, Wales and Northern Ireland 26 November 2021. <https://www.icnarc.org/our-audit/audits/cmp/reports>.
- Xie J, Wu W, Li S, Hu Y, Hu M, Li J, et al. Clinical characteristics and outcomes of critically ill patients with novel coronavirus infectious disease (Covid-19) in China: a retrospective multicenter study. *Intensive Care Med.* 2020;46(10):1863-1872.
- Mokhtari T, Hassani F, Ghaffari N, Ebrahimi B, Yarahmadi A, Hassanzadeh G. Covid-19 and multiorgan failure: A narrative review on potential mechanisms. *J Mol Histol.* 2020;51(6):613-628.
- Covid-19 coronavirus pandemic. <https://www.worldometers.info/coronavirus>. Accessed October 21, 2021.
- Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med.* 2020 May;8(5):475-481.
- Zou X, Li S, Fang M, Hu M, Bian Y, Ling J, et al. Acute Physiology and Chronic Health Evaluation II Score as a Predictor of Hospital Mortality in Patients of Coronavirus Disease 2019. *Crit Care Med.* 2020 Aug;48(8):e657-e665.
- Stephens JR, Stümpfle R, Patel P, Brett S, Broomhead R, Baharlo B, et al. Analysis of Critical Care Severity of Illness Scoring Systems in Patients With Coronavirus Disease 2019: A Retrospective Analysis of Three U.K. ICUs. *Crit Care Med.* 2021 Jan 1;49(1):e105-e107.
- Vandenbrande J, Verbrugge L, Bruckers L, Geebelen L, Geerts E, Callebaut I, et al. Validation of the Acute Physiology and Chronic Health Evaluation (APACHE) II and IV Score in Covid-19 Patients. *Crit Care Res Pract.* 2021 Jun 19;2021:5443083.
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with Covid-19 in Wuhan, China: a retrospective cohort study. *Lancet.* 2020 Mar 28;395(10229):1054-1062.
- Wellbelove Z, Walsh C, Perinpanathan T, Lillie P, Barlow G. Comparing the 4C mortality score for Covid-19 to established scores (CURB65, CRB65, qSOFA, NEWS) for respiratory infection patients *J Infect.* 2021 Mar;82(3):414-451.
- Yildiz H, Castanares-Zapatero D, Hanneke C, Vandermeersch D, Pothén L, Yombi JC. Prospective validation and comparison of COVID-GRAM, NEWS2, 4C mortality score, CURB-65 for the prediction of critical illness in Covid-19 patients. *Infect Dis (Lond).* 2021 Aug;53(8):640-642.
- Bhattacharjee, A, Maji, Rabi L, Gantait, K, Elango, A, Ghosh, S. CT severity score and ABG analysis are the good predictors of mortality in patients with Covid-19 in the tertiary care facility. *IOSR Journal of Dental and Medical Sciences.* 2021;20. 51-57.
- Van Berkel B, Vandevenne J, Coursier K, Alberts V, Van Offenwert J, Verduyck J, et al. Chest CT Diagnosis of Covid-19: Accuracy using CO-RADS and CT-Involvement Scoring. *J Belg Soc Radiol.* 2021 Apr 5;105(1):17.